

- Buijse, A. D., Coops, H., Staras, M., Jans, L., Van Geest, G., Grift, R., . . . Roozen, F. C. (2002). Restoration strategies for river floodplains along large lowland rivers in Europe. *Freshwater Biology*, 47(4), 889-907.
- Erős, T., Kuehne, L., Dolezsai, A., Sommerwerk, N., & Wolter, C. (2019). A systematic review of assessment and conservation management in large floodplain rivers—Actions postponed. *Ecological indicators*, 98, 453-461.
- Grift, R. E., Buijse, A. D., Van Densen, W. L. T., Machiels, M. A. M., Kranenbarg, J., Klein Breteler, J. G. P., & Backx, J. J. G. M. (2003). Suitable habitats for 0-group fish in rehabilitated floodplains along the lower River Rhine. *River Research and Applications*, 19(4), 353-374.
- Palmer, M. A., Hondula, K. L., & Koch, B. J. (2014). Ecological restoration of streams and rivers: shifting strategies and shifting goals. *Annual Review of Ecology, Evolution, and Systematics*, 45, 247-269.
- Pander, J., Mueller, M., & Geist, J. (2018). Habitat diversity and connectivity govern the conservation value of restored aquatic floodplain habitats. *Biological Conservation*, 217, 1-10.
- Stoffers, T., Collas, F. P. L., Buijse, A. D., Geerling, G. W., Jans, L. H., van Kessel, N., . . . Nagelkerke, L. A. J. (2020). 30 years of large river restoration: How long do restored floodplain channels remain suitable for targeted rheophilic fishes in the lower river Rhine? *Science of the Total Environment*, 142931. doi:<https://doi.org/10.1016/j.scitotenv.2020.142931>
- Van Denderen, R. P., Schielen, R. M. J., Westerhof, S. G., Quartel, S., & Hulscher, S. J. M. H. (2019). Explaining artificial side channel dynamics using data analysis and model calculations. *Geomorphology*, 327, 93-110.
- Wolter, C., Buijse, A. D., & Parasiewicz, P. (2016). Temporal and spatial patterns of fish response to hydromorphological processes. *River Research and Applications*, 32(2), 190-201.