

Gebruikte literatuur bij artikel Natte kraamkamers Visionair no. 55, maart 2020

Eick, D. & Thiel, R. (2013). Key environmental variables affecting the ichthyofaunal composition of groynes fields in the middle Elbe River, Germany. *Limnologica* no. 43, p. 297-307.

Gorski, K., De Leeuw, J. J., Winter, H. V., Vekhov, D. A., Minin, A. E., Buijse, A. D., & Nagelkerke, L. A. (2011). Fish recruitment in a large, temperate floodplain: the importance of annual flooding, temperature and habitat complexity. *Freshwater Biology*, 56(11), p. 2210-2225.

Grift, R. E., Buijse, A. D., Van Densen, W. L. T., Machiels, M. A. M., Kranenbarg, J., Klein Breteler, J. G. P., & Backx, J. J. G. M. (2003). Suitable habitats for 0-group fish in rehabilitated floodplains along the lower River Rhine. *River Research and Applications*, 19(4), p. 353-374.

Humphries, P., King, A., McCasker, N., Kopf, R. K., Stoffels, R., Zampatti, B., & Price, A. (2019). Riverscape recruitment: a conceptual synthesis of drivers of fish recruitment in rivers. *Canadian Journal of Fisheries and Aquatic Sciences*, (999), p. 1-13.

Schlösser, I. J. (1991). Stream fish ecology: a landscape perspective. *BioScience*, 41(10), p. 704-712.